


CAZyme™ CthCelH



C5•6 Technologies, Inc.

Technical Specifications

Catalog No. 30555-1 2 mg (0.2 ml)
Lot No.



Store at 4°C. Do not re-freeze.

For *In Vitro* Research Use Only.

Not for Drug or Diagnostic use. Not for use in humans or animals.

Product Description	CAZyme CthCelH, thermostable, recombinant expressed in <i>E. coli</i> cells, cloned from <i>Clostridium thermocellum</i> . 10 mg/ml. MW = 92 kDa
Purity	≥90% pure on Coomassie stained SDS-PAGE.
Recommended Reaction Conditions	CAZyme CthCelH is active between pH 6.0 and 8.0 at 70°C. Optimum pH is 5.8 and optimum temperature is 70°C.
Specific Activity	47 units/mg.
Activity Determination	One cellulase unit will produce 1 micromole of reducing sugar per minute at 70°C from a 1% solution of β-glucan (Megazyme, P-BGBL) in 50 mM sodium acetate at pH 5.8. Assay method available upon request.
Endoglucanase Activities	CAZyme CthCelH possesses <i>endo</i> -cellulase, endoxylanase, arabinoxylanase, β-glucanase and mannanase activities when assayed using insoluble AZCL-linked substrates. Assay method available upon request.
Exoglucanase Activities	CAZyme CthCelH possesses <i>exo</i> -cellulase and cellobiohydrolase activities. Assay method available upon request.
Protein Concentration	10 mg/ml total protein as measured using the Bradford protein assay with BSA as standard.
Stability	Store at 4°C. If properly stored at 4°C, this product is guaranteed for 6 months from date of purchase.
Storage Buffer	50 mM Tris-HCl, pH 7.5, 100 mM NaCl, 25% glycerol.

Note: This enzyme is shipped frozen but should be stored at 4°C. Additional freeze/thaw cycles will result in decreased activity.

C5•6 Technologies 2905 Parmenter St, Middleton, WI 53562 USA
(608) 836.3587
info@c56technologies.com www.c56technologies.com

1

FOR RESEARCH USE ONLY. NOT FOR HUMAN OR DIAGNOSTIC USE

```

MWVGTQPSESAIKSFQELQGRKLDIVHQFINWSTDFSWVRPYADAVYNNNGSILMITWEPW
EYNTVDIKNGKADAYITRMAQDMKAYGKEIWLRLHEANGDWYPWAIGYSSRVNTNETYI
AAFRHIVDIFRANGATNVKWFVNVNCDNVGNGTSYLGHYPGDNYVDYTSIDGYNWGTTS
WGSQWQSFQVFSRAYQALASINKPIIIAEFASAEIGGNKARWITEAYNSIRTSYNKVIA
AVWFHENKETDWRINSSPEALAAAYREAIGAGSSNPTPTPTWTSTPPSSSPKAVDPFEMVR
KMGMGTNLGNTLEAPYEGSWSKSAMEYYFDDFKAAGYKNVRI PVRWDNHTMRTYPYTIDK
AFLDRVEQVVDWLSLRGFVTIINSHHDDWIKEDYNGNIERFEKIWEQIAERFKNKSENLL
FEIMNEPFGNITDEQIDDMNSRILKIIRKTNPTRIVIIGGGYWNSYNTLVNIKIPDDPYL
IGTFHYYPDYEFTHKWRGTWGTQEDMDTVVRVDFVKSWSDRNNIPVYFGEFVMAADR
TSRVKWFYDFISDAALERGFACSVWDNGVFGSLDNDMAIYNRDTRTFDTEILNALFNPPTY
PSYSPKPSPTPRPTKPPVTPAVGEKMLDDFEGVLNWGSYSGEGAKVSTKIIVSGKTGNGME
VSYTGTTDGYWGTVYSLPDGDWSKWLKISFDIKSVDGSANEIRFMIAEKSINGVGDGEHW
VYSITPDSSWKTIEIPFSSFRRLDYQPPGQDMSGTLDLDNIDSIHFMYANNKSGKFVVD
NIKLIGATSDPTPSIKHGHHHHHH

```

Length: 804aa
Theoretical pI: 5.71
Theoretical MW: 91,811 Da
PFAM Structure: GH26 GH5 CBM11
Activity: endo-cellulase
Typical Specific Activity: 47 u/mg
Leader: (-)
Dockerin: (-)
Histag: (+)

Figure 1. Features and sequence of recombinant CAZyme CthCelH (1).




Figure 2. Temperature and pH tolerance of CAZyme CthCelH. Assay conditions available upon request.

1. Yague, E., Beguin, P., and Aubert, J.-P. (1990) Nucleotide sequence and deletion analysis of the cellulase-encoding gene *celH* of *Clostridium thermocellum*. *Gene*. **89**, 61.

C5-6 Technologies 2905 Parmenter St, Middleton, WI 53562 USA
 (608) 836.3587
info@c56technologies.com www.c56technologies.com

